

The effects of the illumination of buildings on house-dwelling bats and its conservation consequences

SÁNDOR BOLDOGH^{1,4}, DÉNES DOBROSI², and PÉTER SAMU³

¹Aggtelek National Park Directorate, Tengersizem-oldal 1, 3758 Jósvafő, Hungary

²Kiskunság National Park Directorate, Liszt Ferenc u. 19, 6000 Kecskemét, Hungary

³Hungarian Bat Research Association, Szabadság u. 13, 5452 Mesterszállás, Hungary

⁴Corresponding author: E-mail: boldogh@anp.kvvm.hu

As the illumination of buildings at night increases, light pollution and negative impacts on wildlife also increase. In order to assess the effect of direct lighting on house-dwelling bats, we examined colonies of *Rhinolophus ferrumequinum*, *Myotis emarginatus* and *M. oxygnathus* in illuminated and non-illuminated buildings found in close proximity to each other. We investigated the onset and timing of nocturnal emergence and measured the body mass and the forearm length of juvenile bats. Results show that bright artificial lighting delays the onset or significantly prolongs the duration of emergence and, in the worst cases, may destroy the whole colony. Juveniles are significantly smaller in illuminated buildings than in non-illuminated ones. The differences in length of the forearm and in body mass may suggest that the parturition time starts later and/or the growth rate is lower in bats living in illuminated buildings. Thus, the illumination of buildings could have serious implications for the conservation of house-dwelling bat colonies.

Key words: light pollution, bat, conservation, artificial roost, *Myotis oxygnathus*, *M. emarginatus*

INTRODUCTION

Light pollution resulting from the increasing illumination of the planet by mankind is having an increasing influence on wildlife (Longcore and Rich, 2004; Rich and Longcore, 2006). The ecological consequences of artificial night lighting are becoming increasingly clear (e.g., Fedum, 1995; Borg, 1996; Harder, 2002; Eisenbeis, 2006; Frank, 2006; Montevecchi, 2006). Scientific data shows that this artificial disturbance also influences nature conservation policy and activities (e.g., Eisenbeis and Hassel, 2000; Health Council, 2000; Le Corre *et al.*, 2002; Salmon 2003, 2006).

Ecological light pollution has obvious effects on bats as well as many other diurnal, crepuscular and nocturnal species. Many groups of insects, of which moths are the most well-known, are attracted in large numbers to lights, and bats are quick to take advantage of these concentrations of prey (Rydell, 1991, 2006; Blake *et al.*, 1994; Rydell and Baagøe, 1996a, 1996b; Gaisler *et al.*, 1998; Swensson and Rydell, 1998). However, artificial illumination not only affects the hunting-ground but may also influence roosts and emergence behaviour (Downs *et al.*, 2003). In 2003, we observed that bats did not emerge after dusk from a church which was directly illuminated by

floodlighting. In another case, a colony disappeared after lights had been installed by the local council. If bats can be restricted or deterred by illumination, this must have important implications for bat conservation. The floodlighting of buildings (mainly churches) was not a common practice in villages in Central and Eastern Europe 10 years ago, but recently it has become increasingly common. The aim of such lighting is to emphasize the attractiveness of the buildings. The abundance of house-dwelling bat colonies in Hungary (see Dobrosi, 1996; Matis *et al.*, 2002; Boldogh, 2006) and the many different practices of lighting make the study particularly relevant.

MATERIALS AND METHODS

Emergence Activity

House-dwelling bat colonies were surveyed in the north and south-east of Hungary (Table 1). The species composition and the size of the roosting colonies were determined beforehand by day.

Firstly, to test whether bright lighting causes any differences in emergence activity, we examined the timing of the nightly onset of emergence and the characteristics of the emergence behaviour in illuminated (roosts 4 and 6) and non-illuminated buildings (roost 7 — Table 1).

Secondly, we disconnected the lights at the illuminated buildings (roosts 4 and 6 — Table 1) for several days (1–3) so the roost-buildings remained dark and examined the onset of nightly emergence and the duration of emergence behaviour. The temporary elimination of lighting was carried out during the days immediately following the basic investigation, in order to avoid and/or reduce the effect of the lunar cycle and the natural shortening of daylength. The investigations were carried out on days with calm weather to avoid the influence of meteorological factors (strong wind, clouds) on the emergence behaviour. During the investigation bats were counted and identified visually and with the help of an ultrasound bat detector (Mini-3 Bat Detector[®]).

The Growth of Juveniles

To investigate the effect of lighting on the growth of juveniles, the length of the forearm and body mass

of bats in the illuminated and the non-illuminated control buildings were compared (Table 1). Control buildings with the same species and similar condition (same type of roof) were selected. The data were collected in the paired colonies on the same day. To minimize disturbance the young bats were usually measured after their mothers had left the roost at dusk; only one parallel measurement was taken in the daytime on warm days (roosts 1 and 4). Random sampling of juveniles was carried out by hand catch. Measurement of pups was carried out immediately after they were captured. The pups of *R. ferrumequinum* were not investigated considering the sensitivity of this species.

We used callipers for measuring the length of the forearm to the nearest 0.1 mm and a spring scale (Pesola Light-Line 50[®]) for measuring body mass. The body mass was measured to the nearest 0.1 g. The analysis of the data was performed with SPSS 12.0[®].

RESULTS

Emergence Activity

Differences in the emergence activity of the bats in the illuminated and non-illuminated buildings were remarkable. Almost all the bats left the undisturbed roosts in the first 30 minutes after dusk, whereas there was a considerable delay in the onset of emergence in the illuminated buildings where most of the bats remained in the roof until the disconnection of the lights (Fig. 1). Some bats flew out but never totally left the site whilst the lights were on. They re-entered repeatedly, flew back into the darkness of the roof and fluttered inside for a long time. The evidence suggests that we can separate the species by their sensitivity: while several *R. ferrumequinum* and *M. oxygnathus* departed when the lights were on, the great majority of *M. emarginatus* remained behind until it was totally dark.

During the first unlit night, the majority of the colony of *M. emarginatus* emerged at the same time as they had done during the previous illuminated nights whilst the

TABLE 1. The details of investigated roosts and colonies

Roost No.	Locality	Roosting species	Date of experiment	Description of illumination	Measured species
1	Kelemér (N Hungary), church	<i>Myotis oxygnathus</i> , <i>M. myotis</i> , <i>Plecotus austriacus</i>	29.06.2006, 16.08.2006	No lights	<i>M. oxygnathus</i> ^{b, c}
2	Alsószuha (N. Hungary), church	<i>Rhinolophus hipposideros</i> , <i>R. ferrumequinum</i> , <i>M. oxygnathus</i> , <i>M. myotis</i> , <i>M. emarginatus</i>	27.06.2006, 16.08.2006	All night ^a	<i>M. oxygnathus</i> ^{b, c}
3	Ragály (N Hungary), church	<i>M. emarginatus</i>	30.06.2005, 27.06.2006	All night from 01.11.2005 ^a	
4	Szólóárdó (N Hungary), church	<i>M. oxygnathus</i> , <i>R. ferrumequinum</i>	29.06.2006, 16.08.2006, 17.08.2006	1 hour (after dusk)	<i>M. oxygnathus</i> ^{b, c, d}
5	Komádi (SE Hungary), castle	<i>M. emarginatus</i>	16.07.2005		<i>M. emarginatus</i> ^{b, c, d}
6	Geszt (SE Hungary), church	<i>R. ferrumequinum</i> , <i>M. emarginatus</i> , <i>Myotis dasycneme</i> , <i>P. austriacus</i>	07–17.07.2003, 15.07.2005	From dusk until 23:30	<i>M. emarginatus</i> ^{b, c, d}
7	Geszt (SE Hungary), castle	<i>R. ferrumequinum</i> , <i>M. emarginatus</i> , <i>P. austriacus</i>	07–17.07.2003	No lights	All species ^d
8	Tatarszentgyörgy (E Hungary), church	<i>M. oxygnathus</i> , <i>Eptesicus serotinus</i>	13.07.2006	From dusk until 23:00	<i>M. oxygnathus</i> ^{b, c}
9	Mezőberény (SE Hungary), church	<i>M. oxygnathus</i> , <i>Nyctalus noctula</i> , <i>P. austriacus</i>	13.07.2006	No lights	<i>M. oxygnathus</i> ^{b, c}

^a – from dusk until dawn

^b – body mass

^c – length of the forearm

^d – emergence activity




FIG. 1. The nightly emergence activity in *M. emarginatus*, in an illuminated (dashed line, roost number 6, $n = 1460$) and a non-illuminated roost (continuous line, roost number 7, $n = 200$) at the same night (07.07.2003). Arrow shows the time when lights were switched off at the illuminated roost

other species (*M. oxygnathus*, *R. ferrumequinum*) flew out earlier. During the second period after the lights had been disconnected, *M. emarginatus* also flew out earlier than they had during the illuminated nights (Fig. 2). *M. emarginatus* was the slowest to adjust to the modified circumstances; *M. oxygnathus* and *R. ferrumequinum* reacted more quickly to the change.

An unfortunate example of the direct effect of illumination was when the largest known *M. emarginatus* colony, consisting of approximately 1,000–1,200 females, left the roost after lights had been installed by

the local council; the floodlights poured light directly through the wide exit-hole and completely flooded the loft (roost 3).

The Growth of Juveniles

The forearm length of juvenile bats was significantly shorter in illuminated than in non-illuminated colonies (Table 2). The difference was greatest during the lactation period (Mann-Whitney *U*-Test, $P < 0.001$) and disappeared by mid-September (Mann-Whitney *U*-Test, $P > 0.05$ — Table 2).




FIG. 2. The nightly emergence activity in *M. oxygnathus* ($n = 230$), during an illuminated (dashed line) (16.07.2006) and the following first non-illuminated night (continuous line) (17.07.2006) at the same roost (roost number 4). Arrow shows the time when lights were switched off during the illuminated nights

The body mass of juveniles was also different between illuminated and non-illuminated colonies. Young bats were larger in dark roosts (Mann-Whitney *U*-Test, $P < 0.001$) and this difference persisted until late summer (Mann-Whitney *U*-Test, $P < 0.015$ — Table 2).

DISCUSSION

Bright artificial illumination negatively affects bats, therefore, the illumination of roosts has serious implications for the conservation of house-dwelling bat colonies. The highest abundance of aerial insects usually occurs around dusk (e.g., Nyholm, 1965; Jones and Rydell, 1994; Rydell *et al.*, 1996); hence bats, especially aerial-hawking bats, emerge from their roosts soon after sunset (e.g., Gaisler, 1963; Herreid and Davis, 1966; Kunz, 1974; Kunz and Anthony, 1996). Our study is consistent with other studies (Erkert, 1982; Kunz and Anthony, 1996) in showing that most bats synchronize the onset of their nightly emergence with sunset. However, we found — correspondingly with Downs *et al.* (2003) — that bright artificial lighting affects the number of bats emerging. Our results even show that lighting delays the onset and the duration of the emergence of bats. Due to such delayed emergence bats miss the highest abundance of aerial insects and lose a significant proportion of their foraging time. In one instance we found that artificial lighting forced the whole colony to leave the roost.

The difference in the length of the forearm may show that the parturition time starts later and/or that the growth rate is slower in bats living in illuminated buildings. Our field observations suggest that both effects occur. The exact time of parturition was not known, therefore, the disparity in age could only be estimated. The length of the forearm can be used for

TABLE 2. The development stages of juvenile bats measured by the length of the forearms (mm) and body mass (g) in the different roosts; *n* — sample size (Mann-Whitney *U*-test)

Species (roost number ^a)	Length of forearm				Body mass				
	Non-illuminated roost		Illuminated roost		Non-illuminated roost		Illuminated roost		<i>P</i> -level
	<i>n</i>	$\bar{x} \pm SD$	<i>n</i>	$\bar{x} \pm SD$	<i>n</i>	$\bar{x} \pm SD$	<i>n</i>	$\bar{x} \pm SD$	
<i>M. emarginatus</i> (5, 6) ^b	53	35.62 ± 4.13	76	31.43 ± 5.14	53	6.07 ± 0.95	76	5.27 ± 1.03	0.001
<i>M. oxygnathus</i> (8, 9) ^b	19	54.95 ± 4.72	23	46.28 ± 7.56	18	14.65 ± 4.65	23	10.74 ± 3.41	0.013
<i>M. oxygnathus</i> (1, 4) ^b	18	46.40 ± 8.31	23	36.46 ± 7.84	19	20.37 ± 2.77	23	14.01 ± 3.16	0.001
<i>M. oxygnathus</i> (1, 2) ^c	18	57.57 ± 1.26	14	56.79 ± 1.63	18	22.96 ± 1.68	14	19.6 ± 1.47	0.001

^a — See Table 1 for roost details and date of measurements

^b — in the lactation period

^c — post-weaning period

estimating the absolute age of juveniles during the rapid and linear phase of growth in the first two weeks (Tuttle and Stevenson, 1982; Anthony, 1988; Kunz and Stern, 1995; Reiter, 2004; Sharifi, 2004a); correct estimation is more difficult for older juveniles (De Paz, 1986; Kunz and Hood, 2000). Since our research involves data from the initial phase, and the gross growth rate in the initial phase is known in *M. oxygnathus* (Sharifi, 2004b) we were able to make a relatively good estimation. However, as Sharifi's (2004b) study was conducted at an undisturbed maternity roost, his results could only be used for estimations at the non-illuminated roosts in our research. Another problem is that there is nothing known about the degree of asynchrony at birth in the different roosts which results in an even more complicated and questionable estimation. Regarding these experiences, we used the detected differences in the length of the forearm only for a rough estimation of disparity in age between the illuminated and non-illuminated colonies. The estimated disparity is at least 7–10(11) days. We had one concrete observation at the early stage of the parturition, when only pregnant females and neonates were found at the illuminated roost while the undisturbed roost had well-developed young. This apparently indicated that birth had been delayed in illuminated buildings.

In natural conditions the growth rate of the forearm in *M. oxygnathus* reaches the plateau about 35–40 days after birth (Sharifi, 2004). Similar rapid progress has also been reported in other free-ranging and captive insectivorous bats in the temperate zone (e.g., Kleiman, 1969; O'Farrell and Studier, 1973; Burnett and Kunz, 1982; Kunz and Anthony, 1982; De Fanis and Jones, 1995; Kunz and Stern, 1995; Swift, 2001; Reiter, 2004). Since we did not find significant differences in the length of forearms between the two different roosts at the end of

summer, it suggests that the individuals at the light-disturbed roosts had also already reached the normal length of the forearm by that time. This equalization may be caused by compensatory growth which has also been reported in other studies (Hoying and Kunz, 1998; Kunz and Hood, 2000). The question of how much the variation is due to different dates of birth and how much to the different rates of growth, is one which deserves further investigation.

In young bats body mass growth rate reflects environmental conditions more responsively than the growth rate of forearms (Kunz and Robson, 1995). The lower availability of insects to lactating females owing to the illumination of roosts, directly leads to lower body mass in juveniles. This is a similar effect to that of bad weather during the maternity period (Kunz and Robson, 1995; Hoying and Kunz, 1998; Kunz and Hood, 2000; Reiter, 2004). The present study shows that in illuminated colonies the body mass remains reduced even after the weaning season. The juvenile bats concerned could probably not compensate for their early disadvantage and/or they had deposited less fat by mid-September. Since hibernation success mainly depends on the body mass achieved, the illumination of maternity roosts may reduce the hibernation success of young bats. Whereas, the earlier born individuals have higher survival rates compared to the ones were born later (Ransome, 1998), the bats born in unlit roosts may have an advantage over those who were born in the illuminated ones.

The conservation strategy for maintaining important house-dwelling colonies is clear: to eliminate direct illumination totally during the whole reproductive season. According to our results, reducing the hours of illumination in the night has little effect: even a one-hour long lighting period after dusk causes significant disruption in behaviour and growth. Summer nights are short in

the temperate zone and even shorter further north, so any artificial reduction in foraging time is disadvantageous.

ACKNOWLEDGEMENTS

We would like to thank Dr. Gábor Csorba, Dr. Zoltán Bihari, Katalin Csilléry and two anonymous reviewers whose constructive comments much improved an earlier version of the manuscript. We are also grateful to János Szilágyi and Ábel Váci for field assistance. Special thanks to Nigel Spring, Gerard Gorman and Fanni B. Szűts who made numerous useful comments and revised the English text. We also have to thank those clerics who let us enter their churches and buildings and made it possible for us to obtain the information for this paper. This work could not have been accomplished without the support of the Aggtelek and Kiskunság National Park Directorates. SB would especially like to say many thanks to his family and to the director of the ANPD Gábor Salamon for providing the creative atmosphere for his work.

LITERATURE CITED

- ANTHONY, E. L. P. 1988. Age determination in bats. Pp. 47–58, *in* Ecological and behavioural methods for the study of bats (T. H. KUNZ, ed.). Smithsonian Institution Press, Washington D.C., 533 pp.
- BLAKE, D., A. M. HUTSON, P. A. RACEY, J. RYDELL, and J. R. SPEAKMAN. 1994. Use of lamplit roads by foraging bats in southern England. *Journal of Zoology (London)*, 234: 453–462.
- BOLDOGH, S. 2006. The bat fauna of the Aggtelek National Park (Hungary) and its surroundings. *Vespertilio*, 9–10: 33–56.
- BORG, V. 1996. Death of the night. *Geographical Magazine*, 68: 56.
- BURNETT, C. D., and T. H. KUNZ. 1982. Growth rates and age estimation in *Eptesicus fuscus* and comparison with *Myotis lucifugus*. *Journal of Mammalogy*, 63: 33–41.
- DE FANIS, E., and G. JONES. 1995. Post-natal growth, mother-infant interactions and development of vocalizations in the vespertilionid bat, *Plecotus auritus*. *Journal of Zoology (London)*, 235: 85–97.
- DE PAZ, O. 1986. Age estimates and postnatal growth in the mouse bat *Myotis myotis* (Borkhausen, 1797) in Guadalajara, Spain. *Mammalia*, 50: 243–251.
- DOBROSI, D. 1996. A handbook for the conservation of bats in Hungary. Hungarian Bat Research Association, Mesterszállás, 48 pp.
- DOWNES, N. C., V. BEATON, J. GUEST, J. POLANSKI, S. L. ROBINSON, and P. A. RACEY. 2003. The effects of illumination the roost entrance on the emergence behaviour of *Pipistrellus pygmaeus*. *Biological Conservation*, 111: 247–252.
- EISENBEIS, G. 2006. Artificial night lighting and insects: attraction of insects to streetlamps in a rural setting in Germany. Pp. 281–304, *in* Ecological consequences of artificial night lighting (C. RICH and T. LONGCORE, eds.). Island Press, Washington D.C., 458 pp.
- EISENBEIS, G., and F. HASSEL. 2000. Attraction of nocturnal insects to street lights: a study of municipal lighting systems in a rural area of Rheinhessen (Germany). *Natur und Landschaft*, 75: 145–156. [in German].
- ERKERT, H. G. 1982. Ecological aspects of bat activity rhythms. Pp. 201–242, *in* Ecology of bats (T. H. KUNZ, ed.). Plenum Press, New York, 425 pp.
- FEDUM, I. 1995. Fatal light attraction. *Journal of Wildlife Rehabilitation*, 18: 10–11.
- FRANK, K. 2006. Effect of artificial night lighting on moths. Pp. 305–344, *in* Ecological consequences of artificial night lighting (C. RICH and T. LONGCORE, eds.). Island Press, Washington D.C., 458 pp.
- GAISLER, J. 1963. Nocturnal activity in the lesser horseshoe bat, *Rhinolophus hipposideros* (Bechstein, 1800). *Zoologické Listy*, 12: 223–230.
- GAISLER, J., J. ZUKAL, Z. REHAK, and M. HOMOLKA. 1998. Habitat preference and flight activity of bats in a city. *Journal of Zoology (London)*, 244: 439–445.
- HARDER, B. 2002. Deprived of darkness: the natural ecology of artificial light at night. *Science News*, 161: 248–249.
- HEALTH COUNCIL. 2000. Impact of outdoor lighting on man and nature. Publication No. 2000/25E, Health Council of the Netherlands, The Hague.
- HERREID, C. F., and R. B. DAVIS. 1966. Flight patterns of bats. *Journal of Mammalogy*, 47: 78–86.
- HOYING, K. M., and T. H. KUNZ. 1998. Variation in size at birth and post-natal growth in the insectivorous bat *Pipistrellus subflavus*. *Journal of Zoology (London)*, 245: 15–27.
- JONES, G., and J. RYDELL. 1994. Foraging strategy and predation risk as factors influencing emergence time in echolocating bats. *Philosophical Transactions of the Royal Society of London*, 346B: 445–455.
- KLEIMAN, D. G., 1969. Maternal care, growth rate and development in the Noctule (*Nyctalus noctula*), pipistrelle (*Pipistrellus pipistrellus*) and serotine

- (*Eptesicus serotinus*) bats. *Journal of Zoology* (London), 157: 187–211.
- KUNZ, T. H. 1974. Feeding ecology of a temperate insectivorous bat (*Myotis velifer*). *Ecology*, 55: 693–771.
- KUNZ, T. H., and E. L. P. ANTHONY. 1982. Age estimation and post-natal growth in the bat *Myotis lucifugus*. *Journal of Mammalogy*, 63: 23–32.
- KUNZ, T. H., and E. L. P. ANTHONY. 1996. Variation in the timing of nightly emergence behaviour in the little brown bat, *Myotis lucifugus* (Chiroptera: Vespertilionidae). Pp. 225–235, in A volume in the memory of J. Knox Jones, Jr. (H. H. GENOWAYS and R. J. BAKER, eds.). *Contributions in Mammalogy*, Texas Tech University, 315 pp.
- KUNZ, T. H., and W. R. HOOD. 2000. Parental care and postnatal growth in the Chiroptera. Pp. 415–454, in *Reproductive biology of bats* (E. CHRICHTON and P. KRUTZSCH, eds.). Academic Press, New York, 510 pp.
- KUNZ, T. H., and S. K. ROBSON. 1995. Postnatal growth and development in the Mexican free-tailed bat (*Tadarida brasiliensis mexicanus*): birth size, growth rates, and age estimation. *Journal of Mammalogy*, 76: 769–783.
- KUNZ, T. H., and A. A. STERN. 1995. Maternal investment and post-natal growth in bats. *Symposium of Zoological Society of London*, 67: 123–138.
- LE CORRE, M., A. OLLIVER, S. RIBES, and P. JOUVENTIN. 2002. Light-induced mortality of petrels: a 4-year study from Réunion Island (Indian Ocean). *Biological Conservation*, 105: 93–102.
- LONGCORE, T., and C. RICH. 2004. Ecological light pollution. *Frontiers in Ecology and the Environment*, 2: 191–198.
- MATIS, Š., P. PIENČÁK, A. KÜRTHY, and E. HAPL. 2002. Overview of summertime findings of bats in the National Park Slovak Karst. *Natura Carpathica*, 43: 195–234. [in Slovakian].
- MONTEVECCHI, W. 2006. Influences of artificial light on marine birds. Pp. 94–113, in *Ecological consequences of artificial night lighting* (C. RICH and T. LONGCORE, eds.). Island Press, Washington D.C., 458 pp.
- NYHOLM, E. 1965. Zur Ökologie von *Myotis mystacinus* (Leisl.) and *M. daubentonii* (Leisl.) (Chiroptera). *Annales Zoologici Fennici*, 2: 77–123.
- O'FARRELL, M. J., and E. H. STUDIER. 1973. Reproduction, growth, and development in *Myotis thysanodes* and *M. lucifugus* (Chiroptera: Vespertilionidae). *Ecology*, 54: 18–30.
- RANSOME, R. D. 1998. The impact of maternity roost conditions on populations of greater horseshoe bats. *English Nature Research Reports*, 292: 1–79.
- REITER, G. 2004. Postnatal growth and reproductive biology of *Rhinolophus hipposideros* (Chiroptera: Rhinolophidae). *Journal of Zoology* (London), 262: 231–241.
- RICH, C., and T. LONGCORE (eds.). 2006. *Ecological consequences of artificial night lighting*. Island Press, Washington, D.C., 458 pp.
- RYDELL, J. 1991. Seasonal use of illuminated areas by foraging northern bats *Eptesicus nilssonii*. *Holarctic Ecology*, 14: 203–207.
- RYDELL, J. 2006. Bats and their insect prey at streetlights. Pp. 43–60, in *Ecological consequences of artificial night lighting* (C. RICH and T. LONGCORE, eds.). Island Press, Washington D.C., 458 pp.
- RYDELL, J., and H. J. BAAGØE. 1996a. Bats and streetlamps. *Bats*, 14: 10–13.
- RYDELL, J., and H. J. BAAGØE. 1996b. Street lamps increase bat predation on moths. *Entomologisk Tidskrift*, 117: 129–135.
- RYDELL, J., A. ENTWISTLE, and P. A. RACEY. 1996. Timing of foraging flights of three species of bats in relation to insect activity and predation risk. *Oikos*, 76: 243–252.
- SALMON, M. 2003. Artificial night lighting and turtles. *Biologist*, 50: 163–168.
- SALMON, M. 2006. Protecting sea turtles from artificial night lighting at Florida's oceanic beach. Pp. 141–168, in *Ecological consequences of artificial night lighting* (C. RICH and T. LONGCORE, eds.). Island Press, Washington, D.C., 458 pp.
- SHARIFI, M. 2004a. Postnatal growth and age estimation in the Mehely's horseshoe bat (*Rhinolophus mehelyi*). *Acta Chiropterologica*, 6: 155–161.
- SHARIFI, M. 2004b. Postnatal growth in *Myotis oxymycterus* (Chiroptera, Vespertilionidae). *Mammalia*, 68: 283–289.
- SVENSSON, A. M., and J. RYDELL. 1998. Mercury vapour lamps interfere with the bat defence of tympanate moths (*Operophtera* spp.; Geometridae). *Animal Behaviour*, 55: 223–226.
- SWIFT, S. M. 2001. Growth rate and development in infant Natterer's bats (*Myotis nattereri*) reared in a flight room. *Acta Chiropterologica*, 3: 217–223.
- TUTTLE, M. D., and D. STEVENSON. 1982. Growth and survival of bats. Pp. 105–150, in *Ecology of bats* (T. H. KUNZ, ed.). Plenum Press, New York, 425 pp.

Received 07 March 2007, accepted 21 May 2007